Keith, E.L. 2013. *Trifolium nigrescens* (Fabaceae), new to the Texas flora. Phytoneuron 2013-32: 1–6. Published 20 May 2013. ISSN 2153 733X

TRIFOLIUM NIGRESCENS (FABACEAE), NEW TO THE TEXAS FLORA

ERIC L. KEITH Raven Environmental Services Inc. P.O. Box 6482 Huntsville, Texas77342 keith@ravenenvironmental.com ek7275@suddenlink.net

Research Associate, Plant Resources Center University of Texas Austin, Texas

ABSTRACT

Trifolium nigrescens is documented as naturalized in Texas. A large population was found growing along a highway right-of-way near Huntsville in Walker County, Texas.

KEY WORDS: Fabaceae, Trifolium, Trifolium nigrescens, Texas, naturalized

A large, naturalized population of *Trifolium nigrescens* Viv. has been discovered in Walker County, Texas. The species has not previously been reported in the state (Correll & Johnston 1970; Hatch et al. 1990; Jones et al. 1997; Turner et al. 2003).

Voucher: **Texas**. Walker Co.: Median of Hwy 30 at intersection with Timberwilde Drive, ca 3 mi W of jct with Interstate 45 on W side of Huntsville, 3 May 2013, *Keith 1034* (TEX, MU).



Figure 1. Large population of Trifolium nigrescens in Walker County, Texas.




Figure 2. Trifolium nigrescens in Walker County, Texas (T. dubium in background).

Trifolium nigrescens is native to southern Europe and southwestern Asia and has been reported as naturalized in the southeastern USA in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, South Carolina, and Tennessee (Isely 1998; BONAP 2011; USDA, NRCS 2013). The Texas population is the westernmost known for the species, and this report increases the number of native and naturalized species of *Trifolium* in the state to sixteen.

The Texas population of Trifolium nigrescens (Figs. 1 and 2), with tens of thousands of plants, was found growing along approximately one kilometer of roadside and median of Hwy 30 west of Huntsville. In Texas, the species most resembles T. repens but can be readily distinguished but its upright rather than prostrate habit (Figs. 1 and 2). It is most similar to T. hybridum, which has not yet been observed in Texas, although it has been recorded near the borders of Texas with Arkansas, Louisiana, New Mexico, and Oklahoma (Isely 1998; BONAP 2011). It can be separated from T. hybridum by its smaller inflorescence of light pink or white flowers (versus pink) and scarious calyx lobe margins and V-shaped sinuses (versus not scarious and broad U-shaped sinuses).

Included below is a key to seventeen species of *Trifolium* most likely to be found in Texas including T. hybridum, which has not yet been documented in the state. The key is derived from Isely (1998). Species distributions follow Diamond et al. (1987), Turner et al. (2003), BONAP (2011), and personal observations.

TRIFOLIUM SPECIES KNOWN AND EXPECTED IN TEXAS

Trifolium amphianthum Torr. & A. Gray

Not Trifolium polymorphum Poir., the South American species, per email communication with Michael Vincent)

Trifolium arvense L.

Trifolium bejarense Moric.

Trifolium campestre Schreber in Sturm

Trifolium carolinianum Michx.

Trifolium dubium Sibthorp

Trifolium hybridum L.

Expected to occcur in Texas

Trifolium incarnatum L.

Trifolium lappaceum L.

Trifolium mucronatum Willd. ex. Spreng.

SYN = Trifolium arizonicum E. Greene

SYN = Trifolium wormskioldii Lehmann var. arizonicum (E. Greene) Barneby

Trifolium nigrescens Viv

Trifolium pratense L.

Trifolium reflexum L.

Trifolium repens L.

Trifolium resupinatum L.

Trifolium subterraneum L.

Trifolium vesiculosum Savi

1. Flowers yellow, common naturalized species.

2. Corolla distinctly striate; inflorescence 0.8–1.3 cm broad with generally 20 or more flowers; 2. Corolla scarcely striate; inflorescence 0.5–0.8 cm broad with 5–20 flowers, petioles of middle

1. Flowers white, pink, purple, red, or bicolored; native or naturalized species.

3. Calyx pubescent.

4. Corolla 4–7 mm long.

Corolla resupinate and bright pink; calyx bladdery inflated in fruit; common naturalized species
 Corolla not resupinate, white or light pink; calyx not inflated in fruit.

7. Inflorescence initially sessile becoming peduncled and bur-like, corolla white turning light pink; calyx lobes stiffly bristly and plumose with stiff trichomes 1–1.5 mm long that are slightly bulbous at base; rapidly spreading naturalized species ... Trifolium lappaceum
7. Inflorescence pedunculate, corolla whitish or lavender commonly turning dull red; calyx lobes lanceolate and subfoliaceous usually with three nerves; common native species ... Trifolium carolinianum

4. Corolla 9–17 mm long.

3. Calyx mostly glabrous (some villous hairs may be present at orifice of tube; wide glabrous but ciliate lobes in *T. bejarense*).

12. Flowers white, only 2–5 outer flowers with petals, the remainder sterile; fruiting heads transformed through curvature of peduncle into a humistrate or subterranean bur; rare naturalized species known only from 2 counties in central Texas Trifolium subterraneum 12. All flowers with petals, flowers white, pink, or red; fruits not as described above.

13. Flowers sessile or subsessile, white, inflorescence large, 2–3 cm wide; calyx tube plainly multistriate with 20 or more nerves; naturalized species widespread in eastern Texas Trifolium vesiculosum 13. Flowers distinctly pedicellate, pedicels ca. 1 mm or more.

14. Inflorescence of two kinds: those that are pedunculate with pink, purple, light red, or bicolored umbellate, petaliferous flowers, and those at ground level, bearing cleistogamous flowers with reduced petals; plants caespitose, substoloniferous or rhizomatous; increasingly rare native species most recently collected 15 years ago Trifolium amphianthum 14. Inflorescence of one kind, bearing white or pink petaliferous flowers.

15. Calyx reticulate-nerved, the lobes > 1 mm in width, ciliate; pedicels villosulous; uncommon native species found in prairies, open post oak woodlands and roadsides generally in the Blackland Prairie and Oak Woods and Prairies Ecoregions Trifolium bejarense

15. Calyx not reticulate-nerved, the lobes < 1 mm, not ciliate; pedicels glabrous.

16. Perennial, caespitose to stoloniferous; peduncles from ground level; calyx lobes < length of the tube; abundant naturalized species found throughout Texas

16. Short lived perennial or annual, caulescent; peduncles not from ground level; calyx lobes < or = tube length; rare naturalized species or not yet recorded in Texas.

17. Inflorescence 1-1.5(-2) cm in diameter; calyx lobes proximally scarious-

ACKNOWLEDGEMENTS

I would like to thank Guy Nesom for reviewing the paper and Michael Vincent for his comments. The population of *Trifolium nigrescens* was discovered in my continued efforts to relocate a living population of the native *Trifolium amphianthum*. A standing reward of "surf and turf" dinner or fine bottle of scotch is offered for the first person that can relocate a living population.

LITERATURE CITED

- Correll, D.S. and M.C. Johnston. 1970. Manual of the Vascular Plants of Texas. Texas Research Foundation, Renner.
- Diamond, D.D., D.H. Riskind, and S.L. Orzell. 1987. A framework for plant community classification in Texas. Texas J. Sci. 39: 203–222.
- Hatch, S.L., K.N. Gandhi, and L.E. Brown. 1990. Checklist of the Vascular Plants of Texas. MP-1655. Texas Agric. Exp. Sta., College Station.
- Isely, D. 1998. Native and Naturalized Leguminosae (Fabaceae) of the United States (Exclusive of Alaska and Hawaii). Monte L. Bean Life Science Museum. Brigham Young Univ., Provo Utah.
- Jones, S.D., J.K. Wipff, and P.M. Montgomery. 1997. Vascular plants of Texas. A comprehensive checklist including synonymy, bibliography, and index. University of Texas Press. Austin.

BONAP. 2011 (last update). North American Plant Atlas (US county-level species maps). Biota of North America Program, Chapel Hill, North Carolina. http://www.bonap.org/genera-list.html
 Turner, B.L., H. Nichols, G. Denny, and O. Doron. 2003. Atlas of the Vascular Plants of Texas. Vol. 2. Sida, Bot. Misc. 24. Botanical Res. Inst. of Texas, Fort Worth.

USDA, NRCS. 2013. The PLANTS Database. National Plant Data Team, Greensboro, North Carolina. Accessed 3 May 2013">http://plants.usda.gov.> Accessed 3 May 2013.

Vincent, M.A. Pers. comm. (3 May 2013). Authority on the genus *Trifolium*. Miami of Ohio Univ., Oxford. Email correspondence.